

Assignment 1: The 1-D Heat Equation (Intro)

Due Tuesday September 14, 2021

John McCuan

August 16, 2021

Problem 1 (*Haberman 1.2.1-3*)

(a) Determine the dimensions of **lineal heat energy density** $\theta = \theta(x, t)$ for which

$$\int_a^b \theta(x, t) dx$$

models the total heat energy in a thin rod found between positions $x = a$ and $x = b$.

(b) Consider a (more complicated) thin rod with varying cross-sectional area $A = A(x)$ and **volumetric heat energy density** also called $\theta = \theta(x, t)$. Determine the expression for the total heat energy between positions $x = a$ and $x = b$.

(c) Derive a heat equation for the temperature $u = u(x, t)$ in the rod of varying cross-sectional area from the previous part under the assumption of constant specific heat capacity c , constant volumetric density ρ , and constant heat conductivity K .

Problem 2 (*Haberman 1.2.8*) Give an expression for the total thermal energy in a rod modeled on an interval $0 \leq x \leq L$ in terms of the temperature $u = u(x, t)$.

Problem 3 If f is a continuous function defined on the interval $[a, b]$ and

$$\int_x^{x+\Delta x} f(\xi) d\xi = 0 \quad \text{whenever } a < x < x + \Delta x < b$$

then explain clearly and carefully why $f((a + b)/2) = 0$.

Problem 4 Find a solution $u = u(x, t)$ of the problem

$$\begin{cases} u_t = u_{xx}, & 0 < x < L \\ u(0, t) = u_0, & t > 0 \\ u(L, t) = u_1, & t > 0 \end{cases}$$

where u_0 and u_1 are given constants.

Problem 5 Find as many solutions $u = u(x, t)$ as you can to the problem

$$\begin{cases} u_t = u_{xx}, & 0 < x < L \\ u_x(0, t) = 1 = u_x(L, t), & t > 0. \end{cases}$$

Do you think you have found all solutions?

Problem 6 Consider functions $u = u(x, t)$ of the form

$$u(x, t) = \frac{1}{L}[u_1(t) - u_0(t)]x + u_0(t).$$

- (a) What can you say about u_0 and u_1 if u is a solution of $u_t = u_{xx}$?
- (b) What if $u_1(t) - u_0(t)$ is a constant but $u_0(t)$ is not a constant?
- (c) Give an example where the conditions in the previous part hold for u_0 and u_1 .

Problem 7 Let u_0 and u_1 be constants. Consider the problem

$$\begin{cases} u_t = u_{xx}, & 0 < x < L \\ u(0, t) = u_0, & t > 0 \\ u(L, t) = u_1, & t > 0 \\ u(x, 0) = g(x), & 0 \leq x \leq L \end{cases}$$

where $g = g(x)$ is a given function. Find the equilibrium solution $u_* = u_*(x)$ associated with this problem and formulate an (interesting) assertion/guess about the relation between the solution $u = u(x, t)$ and the equilibrium solution $u_*(x)$.

Problem 8 (Haberman 1.4.1) Find the equilibrium solution associated with the problem

$$\begin{cases} u_t = u_{xx} + x^2, & 0 < x < L \\ u(0, t) = u_0, & t > 0 \\ u_x(L, t) = 0, & t > 0 \\ u(x, 0) = g(x), & 0 \leq x \leq L. \end{cases}$$

Here u_0 is a given constant and g is a given function.

Problem 9 (Haberman 1.4.2) Consider the one-dimensional heat equation on the interval $0 \leq x \leq L$ with constant conductivity K and internal thermal energy rate-density generation/forcing modeled by $Q(x) = x$.

- (a) Find an expression for the heat energy generated per unit time along the entire rod.
- (b) Find an expression for the rate of heat energy flowing out of the rod at the ends $x = 0$ and at $x = L$.
- (c) What relation should hold between your answers to the first two parts?

Problem 10 (Haberman 1.4.3) Determine the equilibrium temperature distribution for a one-dimensional rod consisting of two different materials in perfect thermal contact at $x = 1$ and satisfying the following conditions:

- (i) The material modeled on $0 \leq x < 1$ has $c\rho = 1$ and $K = 1$ (where c is the specific heat capacity, ρ is the density, and K is the conductivity). Also on $0 \leq x < 1$ there is an internal unit heat source with constant density per time given by $Q = 1$.
- (ii) The material modeled on $1 < x \leq 2$ has $c\rho = 2$ and $K = 2$ and $Q = 0$.
- (iii) $u(0) = 0 = u(2)$.

Perfect thermal contact just means the temperature $u = u(x)$ is continuous at $x = 1$.