

Assignment 14: partial differential equations  
Due Friday, April 28, 2023

John McCuan

May 7, 2023

**Problem 1** (the real gamma function) Consider  $\Gamma : (0, \infty) \rightarrow \mathbb{R}$  by

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt.$$

- (a) Compute  $\Gamma(1)$ .
- (b) Show  $\Gamma(x+1) = x\Gamma(x)$  for  $x > 0$ .
- (c) Show  $\Gamma(n) = (n-1)!$  for  $n = 1, 2, 3, \dots$
- (d) Show  $\Gamma(1/2) = \sqrt{\pi}$ .
- (e) Show

$$\Gamma\left(\frac{1}{2} + n\right) = \frac{(2n)!}{4^n n!} \sqrt{\pi} \quad \text{for } n = 1, 2, 3, \dots$$

- (f) Show the  $n$ -dimensional measure of an  $n$ -dimensional ball of radius  $r$  in  $\mathbb{R}^n$  is  $\omega_n r^n$  where

$$\omega_n = \frac{\pi^{n/2}}{\Gamma(n/2 + 1)}.$$

- (g) Show the  $(n-1)$ -dimensional measure of  $\partial B_r(\mathbf{p})$  is  $n\omega_n$ .
- (h) Specialize the formula from part (f) to the special cases  $n = 2k$  is even and  $n = 2k + 1$  is odd to show  $\omega_n$  is always a rational multiple of a power of  $\pi$ .

**Problem 2** (mean value property) Show that if  $u \in C^2(U)$  is harmonic on an open set  $U \subset \mathbb{R}^n$  and  $B_r(\mathbf{p}) \subset\subset U$ , then

$$u(\mathbf{p}) = \frac{1}{n\omega_n r^{n-1}} \int_{\partial B_r(\mathbf{p})} u.$$

Hint(s): Change variables in the integral so that you integrate over a domain  $B_1(\mathbf{0})$  independent of  $r$ . Differentiate the expression you get with respect to  $r$ , and use the divergence theorem to show the average value is constant. Determine the constant value must be  $u(\mathbf{p})$  (by continuity).

**Problem 3** (heat equation) Find a Fourier sine series/separated variables solution of the heat evolution problem

$$\begin{cases} u_t = \Delta u, & (x, t) \in (0, \pi) \times (0, \infty) \\ u(0, t) = 0 = u(\pi, t), & t > 0 \\ u(x, 0) = \pi/2 - |x - \pi/2|, & 0 \leq x \leq \pi. \end{cases}$$

**Problem 4** (heat equation) Use mathematical software to illustrate the solution you found in Problem 3 above. What is interesting about the (apparent) regularity of the solution?

heat equation with insulated boundary conditions

For Problems 5-7 consider the problem

$$\begin{cases} u_t = \Delta u, & \text{on } (0, \pi) \times (0, \infty) \\ u(0, t) = 0, & t > 0 \\ u_x(\pi, t) = 0, & t > 0 \\ u(x, 0) = x, & 0 \leq x \leq \pi \end{cases} \quad (1)$$

**Problem 5** (heat equation with insulated boundary conditions) Find the **steady state** temperature distribution  $u_0(x)$  for (1).

**Problem 6** (heat equation with insulated boundary conditions) Find a Fourier series/separated variables solution of (1). You will need to find and solve the appropriate Sturm-Liouville problem; you can't just use a sine series.

**Problem 7** (heat equation with insulated boundary conditions) Use mathematical software to illustrate the solution you found in Problem 6 above.

### One-dimensional wave equation on a finite interval

For problems 8 and 9 we consider the following initial/boundary value problem

$$\begin{cases} u_{tt} = u_{xx} & \text{for } (x, t) \in (0, 2) \times [0, \infty) \\ u(x, 0) = x + 1/2 - |x - 1|/2, & x \in [0, 2] \\ u_t(x, 0) = 0, & x \in [0, 2] \\ u(0, t) = 0, & t \geq 0 \\ u(2, t) = 2, & t \geq 0 \end{cases} \quad (2)$$

which is assumed to model the longitudinal deformation of a one-dimensional elastic continuum. I suggest you illustrate the model function  $u$  by representing/plotting a sequence of twenty-one representative parameter/material points  $x_j = j/10$  for  $j = 0, 1, 2, \dots, 20$  as follows: The equilibrium configuration for the elastic continuum is represented by the spatial identity  $u_0(x, t) \equiv x$  as indicated in Figure 1. With




Figure 1: The identity deformation of a one-dimensional continuum

this approach, the initial displacement  $u(x, 0) = x + 1 - |x - 1|$  can be illustrated as indicated in Figure 2 with  $u(1, 0) = 3/2$ .




Figure 2: Initial deformation of a one-dimensional continuum

**Problem 8** (wave equation; Fourier series solution; Boas Chapter 13 Section 4) Let  $w(x, t) = u(x, t) - x$  where  $u$  is the solution of (2), and solve the initial/boundary value problem satisfied by  $w$  with  $w$  given as a superposition of separated variables solutions.

**Problem 9** (wave equation) Animate the function  $u(x, t)$  obtained in the previous problem using the mapping approach illustrated in Figures 1 and 2 above (with time  $t$  as the animation parameter).

**Problem 10** (integral identities for the multi-dimensional wave equation) Consider the initial/boundary value problem

$$\begin{cases} u_{tt} = \Delta u & \text{on } U \times (0, \infty) \\ u(\mathbf{x}, 0) = u_0(\mathbf{x}), & \mathbf{x} \in U \\ u_t(\mathbf{x}, 0) = v_0(\mathbf{x}), & \mathbf{x} \in U \\ u(\mathbf{x}, t) = \phi(\mathbf{x}, t), & \text{on } \partial U \times [0, \infty) \end{cases} \quad (3)$$

where  $U$  is a bounded domain with  $C^1$  boundary in  $\mathbb{R}^n$  and  $u_0, v_0$ , and  $\phi$  are given smooth functions. Assume  $u \in C^2(\bar{U} \times [0, \infty))$  is a solution of (3) and consider the “energy” quantity

$$E(t) = \frac{1}{2} \int_U [u_t^2 + |Du|^2] \quad (4)$$

which may be considered as a sum of kinetic and potential energies.

(a) Calculate the derivative

$$\frac{dE}{dt}$$

and use the divergence theorem to express this quantity in terms of the boundary values.

(b) Give conditions on the function  $\phi$  under which the energy is conserved, i.e.,

$$\frac{dE}{dt} \equiv 0.$$

(c) Interpret the conditions you gave in part (b) above in terms of physical model assumptions for a two-dimensional ( $n = 2$ ) elastic membrane.

### Bonus Problems

**Problem 11** (calculus of variations) Formulate/model the energy (potential energy due to gravity) associated with a symmetric hanging chain given as the graph of a function  $u \in C^1[-1, 1]$  with  $u(-1) = 0 = u(1)$  and length  $L = 4$ .

**Problem 12** (calculus of variations) Use the method of Lagrange multipliers to find an ODE satisfied by the model function  $u$  of the previous problem:

(a) Let

$$\mathcal{B} = \left\{ u \in C^1[-1, 1] : u(-1) = 0 = u(1), \text{ and } \int_{-1}^1 \sqrt{1 + u'(x)^2} dx = 4 \right\}$$

Consider  $\mathcal{F}[u] = \mathcal{E}[u] - \lambda \text{length}[u]$  where  $\mathcal{E}$  is the potential energy. Show that if  $u_0 \in C^2[-1, 1] \cap \mathcal{B}$  satisfies

$$\mathcal{E}[u_0] \leq \mathcal{E}[u] \quad \text{for all } u \in \mathcal{B},$$

then there exists some  $\lambda \in \mathbb{R}$  such that  $\mathcal{F}[u_0] \leq \mathcal{F}[u]$  for all

$$u \in \mathcal{A} = \{w \in C^1[-1, 1] : w(-1) = 0 = w(1)\}.$$

(b) Compute the first variation  $\delta\mathcal{F}_{u_0}[\phi]$  for  $\phi \in C_c^\infty[-1, 1]$ , and use the fundamental lemma of the calculus of variations to find all  $C^2$  minimizers  $u_0$  of  $\mathcal{F}$ .

(c) Solve the ODE from part (b) above, and numerically find  $\lambda$  to find the model shape of the hanging chain. Hint:

$$\frac{d}{dx} \cosh^{-1}(x) = \frac{1}{\sqrt{x^2 - 1}}.$$

**Problem 13** Use the potential energy due to gravity along with the elasticity model for the tension force in the hanging slinky to model the hanging slinky using the calculus of variations.